



Chief Executive: Joe Harrison

Chairman: Simon Lloyd

Request under Freedom of Information Act 2000

Thank you for your request for information which we received on 10 May 2021.

I am pleased to confirm the following.

- 1. How many serious incidents occurred during surgery in the years 2018, 2019, 2020, and 2021 up to the date received (07/05/2021)? Please provide a breakdown by year.
 - For each serious incident, please provide as much detail as possible within the time/cost framework, and without any identifying information that could incur a personal data exemption.
 - Please provide a breakdown by i) never events ii) all other serious incidents

care of the deteriorating patient Description	Date Reported	SUI Categories	Description	Location	Severity	Never Event?
DATE: 12 with a marginery blood commenced. SHIELD CENTER SHIEL ALL SHIELD CENTER SHIEL ALL SHIELD CENTER SHIEL ALL SHIELD CENTER SHIEL ALL SHIELD CENTER SHIELD CENTE	15-03-18	care of the	03:34: Baby delivered.		Moderate	No
Dec 12 - 724 MOH Call as support staff had not yet attended from first MOH call for biblood products, faborur was decentions a state of the control of t		_				
to fiblood products, labour ward conditation attended on the production of the consultant. General surgeons present in the attraction of the consultant. General surgeons present in the attraction of the consultant. General surgeons present in the attraction of the consultant. General surgeons present in the attraction of the consultant and the consultan						
concultant seven on the hatter. Blooding had settled. (SSEC concultant arrived in heater, blooding had settled. (Total SM 43540) (Secondly a tooth which were dued in the MOT of left ultimic it was noticed that two the cather which were due to be extracted were still present. Secondly a tooth which was not part of the extraction plan had been removed. (Concerns over recent weeks about the rate of revision following primary hip replacement at MOCUIT the 5 – and 30-year revision rates as described by the National Joint Registry are higher than we would expect, and the issue seems to have became non-promise in this bat 2-years. Concerns about a number of potential infections (Infection is associated with revision) observed by constitute of the high seven and the second of the promise in the MOT of the year is in part because of incleaded virginarie due to 12 above. A background of 37/4584 deep infections (1.9%) in the 3s months to september 2039, following hip or line or episcement surgery. This is not a statistical outlier ITCD Lab higher than the historical populorin TDC. Death (general) At 88 20 aboptimal care of the deteriorating patient the deteriorating patient of the deteriorating patient the deterioration patient the deterioration patient the deteriorating patient the deterioration patient the deterioration patient			, ,			
05.00 Consolation arrived in theatire, beeding had settled. Total Ed. 454ml.						
Apatient was reversed in the MID cleft clinic. It was noticed that two the control with over due to be extracted were still present.			05:00: Consultant arrived in theatre, bleeding had settled.			
Lestin which were due to be extracted were still present. Secondly a tooth which was not part of the extraction plan had been remixed. Concerns over recent weeks about the rate of revision following primary hip replacement at MRUII: the 5 - and 10 year revision rates as described by the National laint Regestry and specif and were the time to the National laint Regestry and specif and were the time that 2 syeas. Concerns about a number of potential infection (infection is associated with revision) observed by consultant orthopsedic surgeons over recent weeks. It is not yet clear whether these cases will be confirmed microbiologically as deep base infection and / or if they state in part because of interesed vigilization for 10 states. A background of 17/853 deep infections (1.9%) in the 1s months to September 2019, following fip or time replacement surgery. This is not a statistical outler frido but is higher than the historical position (TBC). A background of 17/853 deep infections (1.9%) in the 1s months to September 2019, following prior or replacement surgery. This is not a statistical outler frido but is higher than the historical position (TBC). A background of 17/853 deep infections (1.9%) in the 1s months to September 2019, following prior or prior the state of the deteriorating patient. A background of 17/853 deep infections (1.9%) in the 1s months to September 2019, following prior or prior the state of the 1state of t	08-06-18	Surgical Error		Other	Moderate	Yes
Discrete to the content of the conte		3	1 '			
replacement at MKURF the 5 - and 10 year revision rates as described by the National Joint Registry are higher than would expect, and the issue seems to have become more prominent in the last 2-3 years. Concerns about a number of potential infections (infection is associated with revision) observed by consultant orthospaedic surgipions over recent weeks, it does not seem that the revision observed by consultant orthospaedic surgipions over recent weeks, it does not seem that the revision observed by consultant orthospaedic surgipions over recent weeks, it does not seem that the revision of 12 (2655 deep infections (115%) in the 15 months to September 2025, following high or knee replacement surgery. This is not a statistical outlier (1802 but is higher than the historical position (1802). 30.10-8.20 Sub-optimal Pather had 3480ml Bill, following before the historical position (1802). Pather had 3480ml Bill, following before the historical position (1802). Pather had 3480ml Bill, following before the historical position (1802). Pather had 3480ml Bill, following before that the provious ELCS). Pather had 3480ml Bill, following before the sub-optimization to go to the hater. In the deteriorating pather than the sub-optimization of the company of the deteriorating pather than the sub-optimization of the company of the sub-optimization to go to the sub-optimization of the company of the sub-optimization to the company of the sub-optimization of the sub-optimization of the company of the sub-optimization of the sub-optimizati						
National Joint Registry are higher than we would expect, and the issue seems to have become more premiment in the last 2-3 years. Concerns about a number of potential infections (Infection is associated with revision) observed by consultant orthopaedic surgeons over recent weeks. It is notly to clear whether these cases will be confirmed introllogically as deep tissue infection and / or if they arise in part because of increased vigilance due to (1) above. A background of 12/635 deep infections (19%) in the 15 months to September 2019, following hip or knee replacement surgery. This is not a statistical outlier (TRC) but is higher than the historical position (TRC). Patient had 1488mt EBL following uterine ruputure. Patient had 1488mt EBL following uterine ruputure. Patient had 1488mt EBL following uterine ruputure. Patient had 1488mt EBL following the following bedside US abdominals (and phaemobate deteriorating patient care of the deteriorating patient in the patient of the	06-12-19	Unit Closure			Moderate	No
Concerns about a number of potential infections (infection is associated with revision) observed by consultant orthopaedic surgeons over recent weeks. It is not yet dear whether these cases will be confirmed mitoriologically as deep tissue infection and / or if they arise in part because of increased vigliance due to (i) above. A background of 12/635 deep infections (1,9%) in the 15 months to september 2019, following thip or here replacement surgery. This is not a statistical outlier (TBC) but is higher than the historical position (TBC). Patient had 1488ml Bill following tenine ruputure. Phase 1 Patient had 1488ml Bill following tenine ruputure. A background of 12/635 deep infections (1,9%) in the 15 months to a statistical outlier (TBC) but is higher than the historical position (TBC). Patient had 1488ml Bill following tenine ruputure. Phase 1 Patient had 1488ml Bill following tenine ruputure. A background of 12/635 deep infections (1,9%) in the 15 months to the 15 months of the 15			National Joint Registry are higher than we would expect, and the issue seems			
revision) observed by consultant orthopaedic surgeons over recent weeks. It is not yet clear whether these cases will be confirmed incrobiologically as deep itssue infection and / or if they arise in part because of increased vigilance due to [1] above. A background of 12/635 deep infections (1.9%) in the 15 months to September 2019, following hip or kneer replacement surgery. This is not a statistical outler (TISU) but is higher than the historical position (TIBC). Sub-optimal Patient had 1886 ml BLL following uterine ruppture. Managed well with Chyototin / 40/U/ transcamic add/ haemobate deteriorating patient 14-08-20			, ,			
deep issue infection and / or if they arise in part because of increased vigilance due to (a) above. A background of 12/633 deep infections (1.9%) in the 15 months to September 2019, following hip or kneer replacement surgery. This is not a statistical outler (1805) but is higher than the historical position (1805). 20-08-20 Sub-optimal care of the Managed well with Oxyotorial / 401// Tranexamic acid/ haemobate Phase 1 20-08-20 Sub-optimal patient Patient had 1488ml EBL following uterine ruputure. Phase 1 20-08-20 Sub-optimal patient Patient had 1488ml EBL following uterine ruputure. Phase 1 20-08-20 Sub-optimal patient Patient had 1488ml EBL following uterine ruputure. Phase 1 20-08-20 Sub-optimal patient Patient had 1488ml EBL following bedside Use 3 baddenial scan to go to theatre. Phase 1 20-08-20 Sub-optimal patient Patient had 20-09 patient Phase 1 20-09-20 Sub-optimal patient Patient had 20-09 patient Patient admitted from ED for emergency laparoscopic appendicactomy. Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Patient Admitted from ED for emergency laparoscopic appendicactomy. Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Patient Had 1800 patient Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Patient Had 1800 patient Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Phase 1 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Error Patient Had 1800 patient Phase 2 20-09-20 Surgical Err			,			
Vigilance due to (1) above.			· · · · · · · · · · · · · · · · · · ·			
September 2015, following hij or kneer replacement surgery. This is not a statistical outlier (TBC) but is higher than the historical position (TBC). 201-08-20 Sub-optimal care of the deteriorating patient care of the control of the deteriorating patient in the deterioration performed. Past induction whilst in the Anaesthetic room, SATS dropped into the 80s, then into the 70s. Patient then had a cardiac arrests. 19-10-20 Surgical Error Patient had Right Color dyratial kneer replacement in the morning, I was the scrib parcititioner for the procedure. The procedure went smoothly, when component saind both implants were opened to the scrub practitioner and inbilation ponents and both implants were opened to the scrub practitioner and inbilation ponents and both implants were opened to the scrub practitioner and implanted by the surgeon detected the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to the determinance of the process of the determinance of the determinance of the determ						
September 2015, following hij or kneer replacement surgery. This is not a statistical outlier (TBC) but is higher than the historical position (TBC). 201-08-20 Sub-optimal care of the deteriorating patient care of the control of the deteriorating patient in the deterioration performed. Past induction whilst in the Anaesthetic room, SATS dropped into the 80s, then into the 70s. Patient then had a cardiac arrests. 19-10-20 Surgical Error Patient had Right Color dyratial kneer replacement in the morning, I was the scrib parcititioner for the procedure. The procedure went smoothly, when component saind both implants were opened to the scrub practitioner and inbilation ponents and both implants were opened to the scrub practitioner and inbilation ponents and both implants were opened to the scrub practitioner and implanted by the surgeon detected the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to the determinance of the process of the determinance of the determinance of the determ			A background of 12/635 deep infections (1.9%) in the 15 months to			
care of the deteriorating patient Sub-optimal care of the deteriorating patient Sub-optimal care of the deteriorating by the sub-optimal care of the deteriorating patient Sub-optimal care of the deteriorating by the sub-optimal care of the deteriorating patient may be sub-optimal care of the deterioration pour laters he lost a further 700ml while in transit to labour ward. decision was made following bedside US abdominal scan to go to theatre. In theatre a further 1.6L was lost. Lotal ESL was 31. Unexpected Death (general) Death (general) Ansesthetic induction performed. Phase 1 mit beatre a further 1.6L was lost. Lotal ESL was 31. Unexpected Death (general) Patient and Right form ED for mereginery laparoscopic appendicectomy. The surface of the procedure of the surface patient and state of the procedure of the surface patient and state of the procedure of the surface patient and state of the procedure went smoothly, when component staining was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and saked the circulating staff to open the implant affect hecking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 37:15 hours, I am submitting the above procedure to NIR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NIR will lateful friber to be implant incompatibility but in this occasion it did not alert. Unexpected admission to NICU Maternity Emergency LSCS for fetal distress, absent fetal movements. Daby born in poor condition Arterial PH - 6.92 BE -3.72 Lac - 13.3 Venous PH - 7.17 BE -7						
care of the deteriorating patient Sub-optimal care of the deteriorating patient Sub-optimal care of the deteriorating by the sub-optimal care of the deteriorating patient Sub-optimal care of the deteriorating by the sub-optimal care of the deteriorating patient may be sub-optimal care of the deterioration pour laters he lost a further 700ml while in transit to labour ward. decision was made following bedside US abdominal scan to go to theatre. In theatre a further 1.6L was lost. Lotal ESL was 31. Unexpected Death (general) Death (general) Ansesthetic induction performed. Phase 1 mit beatre a further 1.6L was lost. Lotal ESL was 31. Unexpected Death (general) Patient and Right form ED for mereginery laparoscopic appendicectomy. The surface of the procedure of the surface patient and state of the procedure of the surface patient and state of the procedure of the surface patient and state of the procedure went smoothly, when component staining was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and saked the circulating staff to open the implant affect hecking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 37:15 hours, I am submitting the above procedure to NIR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NIR will lateful friber to be implant incompatibility but in this occasion it did not alert. Unexpected admission to NICU Maternity Emergency LSCS for fetal distress, absent fetal movements. Daby born in poor condition Arterial PH - 6.92 BE -3.72 Lac - 13.3 Venous PH - 7.17 BE -7	01-08-20	Sub-optimal	Patient had 1488ml EBL following uterine ruputure.	Theatres -	Moderate	No
patient descriptions of the description of the desc		care of the	· · · · · · · · · · · · · · · · · · ·			-
Sub-optimal care of the deteriorating patient with the patient precovery and 100ml in ward 90 and admission. hours later she lost a further 700ml while in transit to labour ward, decision was made following bedieful Sabdominal scan to go to theatre. In theatre a further 1.6L was lost. total EBL was 31. 19-10-20 Unexpected Death (general) Patient admitted from ED for emergency laparoscopic appendicectomy. Anaesthetic induction performs with the into the 70-8. Patient then had a cardiac arrest. Patient had Right Oxford partial knee replacement in the morning. I was the into the 70-8. Patient then had a cardiac arrest. 19-10-20 Surgical Error Patient had Right Oxford partial knee replacement in the morning. I was the component sizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tiblal components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17-15 hours, I am submitting the above procedure to NIR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NIR will alert if there is an implant incompatibility but in this occasion it did not alert. Unexpected admission to NICU Maternity Service - Baby delivered. Significant mec at delivery. Taken to resus. Apgars 4 (1),5 (5),6 (10). Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. Baby delivered and and patient was the patient was the patient was the patient was and oxygen required. Apgars 4 (1),5 (5),6 (10). Apgars 4 (1),5 (5),6 (10). Arterial PH - 6.92 BE - 13.72 Lea - 13.13 Venous PH - 7.17 BE - 7.24		_				
deteriorating patient bours later she lost a further 700ml while in transit to labour ward, decision was made following bedside US abdominal scan to go to theatre. In theatre a further 1.6. was lost, total EBL was 3L. Patient admitted from ED for emergency laparoscopic appendicectomy. Patient admitted from ED for emergency laparoscopic appendicectomy. Phase 1 Patient admitted from ED for emergency laparoscopic appendicectomy. Phase 1 Post induction which in the Anaesthetic room, SATS dropped into the 80s, then into the 70s. Patient then had a cardiac arrest. Patient had Right Dxford partial knee replacement in the morning. I was the scrub practitioner for the procedure. The procedure went smoothly, when component sizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the menical bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17-15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will allert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to resus. Baby delivered. Significant mec at delivery. Taken to resus. Appars 4 (1),5 (5),6 (10) Transfer to NNU Moderate Phase 1 Maternity Exercice - Unexpected admission to NICU Maternity Exercice - Unexp	24-08-20	Sub-optimal			Moderate	No
In theatre a further 1.6. was lost, total EBL was 3.1 Patent admitted from ED for emergency laparoscopic appendicectomy. Patent admitted from ED for emergency laparoscopic appendicectomy. Phase 1 Post induction white from ED for emergency laparoscopic appendicectomy. Phase 1 Post induction white in the Anaesthetic room, SATS dropped into the 80s, then into the 70s. Patient then had a cardiac arrest. Patient had Right Oxford partial knee replacement in the morning. I was the scrub practitioner for the procedure. The procedure went smoothly, when component sizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibla components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the menical bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to result in the result of the service of				Luase 1		
Death (general) Patient admitted from ED for emergency laparoscopic appendicectomy. Phase 1 Phase 2 Phase 2 Phase 2 Phase 3 Phase 3 Phase 4 Phase 2 Phase 2 Phase 3 Phase 4 Phase 4 Phase 2 Phase 2 Phase 3 Phase 4 Phase 4 Phase 2 Phase 4 Phase 2 Phase 2 Phase 3 Phase 4 Phase 4 Phase 4 Phase 5 Phase 5 Phase 6 Phase 7 Phase 8 Phase 8 Phase 9 Phase 1 Phase 9 Phase 1 Phase		patient				
Post induction whilst in the Anaesthetic room, SATS dropped into the 80s, then into the 70s. Patient then had a cardiac arrest. Patient had Right Oxford partial knee replacement in the morning. I was the scrub practitioner for the procedure. The procedure went smoothly, when components slizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NIR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NIR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to resus. NICU Maternity Service - Unexpected admission to NICU Maternity Service - Unexpected admission to NICU Maternity Service - Unexpected admission to NICU After and pead Reg attended inflation breaths and oxygen required. Agars 4 (1),5 (5),6 (10) Transfer to NNU After in port condition cord around the neck 3 times and true knot in cord. aggrs 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Afterial PH - 6.92 BE - 1.3.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24	28-08-20		Patient admitted from ED for emergency laparoscopic appendicectomy.		Major	No
19-10-20 Surgical Error Patient had Right Oxford partial knee replacement in the morning. I was the scrub practitioner for the procedure. The procedure went smoothly, when component sizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Baby delivered. Significant mec at delivery. Baby delivered. Significant mec at delivery. Taken to resus. NICU Read present. Taken to resus. NiCU Read present. Taken to resus. Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. Baby delivered. Significant mec at delivery. Service - Unexpected admission to NICU Emergency LSCS for fetal distress, absent fetal movements. Baby delivered. Significant mec at the full movements. Baby byto ni no por condition cord around the neck 3 times and true knot in cord. apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE - 13.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24 Architecture of the procedure is the scrub and procedure is the procedure in the scrub full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE - 7.24 Architecture is the procedure is the		Death (general)	Post induction whilst in the Anaesthetic room, SATS dropped into the 80s,	Phase 1		
component sizing was done, the circulating personnel checked and confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to resus. Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU Apgars 0 (1),5 (1),6 (10) Transfer to NNU Apgars 0 (2) Imin / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE - 13.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24	19-10-20	Surgical Error	†		Moderate	Yes
confirmed with the operating surgeon the femoral and tibial components and both implants were opened to the scrub practitioner and implanted by the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alter it fithere is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Unexpected admission to NICU NiCU Baby delivered. Significant mec at delivery. Taken to resus. Neonatal emergency call made. Senior midwife and pead Reg attended inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Bergency LSCS for fetal distress, absent fetal movements. baby born in poor condition cord around the neck 3 times and true knot in cord. apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 La - 11.3 Venous PH - 7.17 BE - 7.24				Phase 2		
the surgeon. The surgeon checked the thickness of the meniscal bearing required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to resus. Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU Maternity Service - Unexpected cord around the neck 3 times and true knot in cord. apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE - 13.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24			confirmed with the operating surgeon the femoral and tibial components			
required and asked the circulating staff to open the implant after checking it. The bearing was inserted, wound closed and patient was transferred to recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Baby delivered. Significant mec at delivery. Taken to resus. NICU Neonatal emergency call made. Senior midwife and pead Reg attended inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Transfer to NNU Maternity Service - Unexpected admission to NICU NICU Maternity Service - Unexpected admission to NICU Apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE - 13.72 Lac - 11.3 Venus PH - 7.17 BE - 7.24						
recovery. Around 17:15 hours, I am submitting the above procedure to NJR when I noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. D4-01-21			required and asked the circulating staff to open the implant after checking it.			
noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Unexpected admission to NICU NICU NICU Neonatal emergency call made. Senior midwife and pead Reg attended inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. Service - Unexpected admission to NICU NICU NICU NICU NICU NICU Maternity Service - Unexpected admission to NICU Aggars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24						
noticed that the meniscal bearing inserted was labelled for the left side. Usually the NJR will alert if there is an implant incompatibility but in this occasion it did not alert. Decision made to go for CAT 2 EMCS due to suspicious CTG. Pead present. Unexpected admission to NICU NICU NICU Neonatal emergency call made. Senior midwife and pead Reg attended inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. Service - Unexpected admission to NICU NICU NICU NICU NICU NICU Maternity Service - Unexpected admission to NICU Aggars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24			Around 17:15 hours, I am submitting the above procedure to NIR when I			
Decision made to go for CAT 2 EMCS due to suspicious CTG. Service - Unexpected admission to NICU Maternity Service - Unexpected admission to NICU Maternity Service - Unexpected admission to NICU Maternity Service - Unexpected Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. Service - Unexpected admission to NICU Maternity Service - Unexpected admission to NICU Apgars 9 (1) min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24			noticed that the meniscal bearing inserted was labelled for the left side.			
Service - Unexpected admission to NICU Apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24	04-01-21	Maternity		Theatres	Moderato	No
admission to NICU Taken to resus. Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24	04-01-21	Service -	Pead present.		Moderate	.,,0
NICU Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU Service - Unexpected admission to NICU Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24 Neonatal emergency call made. Senior midwife and pead Reg attended Inflation breaths and pead Reg attended Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Theatres - Phase 1 No Phase 1 Arterial PH - 6.92 BF -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24						
Inflation breaths and oxygen required. Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU BE - 13.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24			Neonatal emergency call made.			
Apgars 4 (1),5 (5),6 (10) Transfer to NNU Maternity Service - Unexpected admission to NICU Apgars 9 (2),5 (5),6 (10) Transfer to NNU Emergency LSCS for fetal distress, absent fetal movements. baby born in poor condition Cord around the neck 3 times and true knot in cord. Apgars 9 (2) 1 min / 3 (2) 5 mins / 4 (2) 1 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24						
Maternity Service - Unexpected admission to NICU Bright Ph - 6.92 BE - 13.72 Lac - 11.3 Venous PH - 7.17 BE - 7.24 Major Phase 1 Theatres - Phase 1 Theatres - Phase 1 Theatres - Phase 1 Theatres - Phase 1 And in the phase 1 Theatres - Phase 1 Theatres - Phase 1 Theatres - Phase 1 And in theatres - Phase 1 Value - Phase 1 Theatres - Phase 1 Theatres - Phase 1 Theatres - Phase 1 Value - Phase 1 Venous Phase 1 Value - Phase 1			Apgars 4 (1),5 (5),6 (10)			
Service - baby born in poor condition cord around the neck 3 times and true knot in cord. NICU apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24	20.04.5:					
Unexpected admission to NICU apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24	30-04-21				Major	No
NICU apgars 0 @ 1 min / 3 @ 5 mins / 4 @ 10 minutes full neonatal resuscitation and intubation including cardiac compression Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24		Unexpected admission to NICU				
Arterial PH - 6.92 BE -13.72 Lac - 11.3 Venous PH - 7.17 BE -7.24						
Lac - 11.3 Venous PH - 7.17 BE -7.24						
Venous PH - 7.17 BE -7.24						
baby admitted to NNU and to be transferred out for cooling.			baby admitted to NNU and to be transferred out for cooling.			

You are advised that this information is provided in accordance with the Freedom of Information Act 2000 and is for your personal use. Any re-use of this information will be subject to copyright and the Re-Use of Public Sector Information Regulations (1st July 05) and authorisation from Milton Keynes Hospital NHS Foundation Trust will be required. In the event of any re-use, the information must be reproduced accurately and not used in a misleading manner.

If you are unhappy with the information received in response to this request, please address your complaint to the Patient Affairs Office at Milton Keynes Hospital NHS Foundation Trust, Standing Way, Eaglestone, Milton Keynes MK6 5LD If, after exhausting our internal process, you are still unhappy with the information received, you may write to the Information Commissioner's Office, Wycliffe House, Water Lane, Wilmslow, Cheshire SK9 5AF.

If you need any further assistance, please do not hesitate to contact us at the address above.

Yours sincerely,

Freedom of Information Co-ordinator For and on behalf of Milton Keynes Hospital NHS Foundation Trust

Any re-use of this information will be subject to the 'Re-use of Public Sector Information Regulations' and best practice.