

## Distal Radius Fracture in Adults Questionnaire

1. Which hospital do you work at?

Milton keynes University Hospital
-----------------------------------

2. Does your hospital accept or manage trauma patients? 

Yes	x	No	
-----	---	----	--

3. Is your hospital a designated major trauma centre? 

Yes		No	x
-----	--	----	---

4. How many adults with closed distal radius fracture does your hospital manage in a month?

~50
-----

5. Does your department have a written guideline for the investigation and management of potential closed distal radius fracture following trauma?

Yes	
No	
Verbal Only	x

6. If yes, where is your guideline taken from? (For example, BOAST, locally derived guideline etc.)

--

7. For patients requiring a closed manipulation of their fracture that present *during the day*, where is this manipulation carried out?

Plaster room	
In the emergency department (excluding resuscitation area)	x
Resuscitation bay in the emergency department	x
Operating Theatre	x
Other (please specify)	

Please specify:

8. Which specialty is responsible for the initial manipulation of the fracture?

Emergency Medicine	x	Trauma and Orthopaedics	x	Other (please specify)	
--------------------	---	-------------------------	---	------------------------	--

9. What form of analgesia is most commonly used for the manipulation procedure?

Regional nerve blockade	
Regional intravenous anaesthesia (Bier's block)	
Local haematoma block	x
Procedural sedation and analgesia (please specify)	x
General anaesthetic	x

Please specify:

No specific method of analgesia specified	
---	--

10. If a Bier's block is performed, which specialty performs the Bier's block?

Emergency Medicine		Anaesthetics	
Trauma and Orthopaedics		Other (please specify)	

Please specify:

11. What grade of doctor is most commonly responsible for performing the Bier's Block?

--

12. For patients requiring a closed manipulation of their fracture that present *overnight*, where is this manipulation carried out?

Plaster room	
In the emergency department (excluding resuscitation area)	x
Resuscitation bay in the emergency department	x
Operating Theatre	x
Other (please specify)	
Manipulation not carried out overnight	

Please specify:

13. Overnight, which specialty is responsible for the initial manipulation of the fracture?

Emergency Medicine	x	Manipulation not carried out overnight	
Trauma and Orthopaedics		Other (please specify)	

Please specify:

14. Overnight, what form of analgesia is used for the manipulation procedure?

Regional nerve blockade	
Regional intravenous anaesthesia (Bier's block)	
Local haematoma block	x
Procedural sedation (please specify)	x
General anaesthetic	x
No specific method of analgesia specified	
Manipulation not carried out overnight	

Please specify:

15. If patients do not receive manipulation out-of-hours where and when do they return?

<i>Where</i>		<i>When</i>	
Emergency Department	x	Following morning	x
Fracture clinic		Next working day (i.e. on Monday if seen over the weekend)	
Plaster room		Next available routine fracture clinic	

Within 24 hours	
-----------------	--

16. Following presentation to hospital with a distal fracture, how soon after the injury are patients the Fracture Clinic?

Within 48 hours	x
Within 72 hours	
More than 3 days later	

radius  
seen in